


# Andorra

Total population: 78 000 > Population aged 15 years and older (15+): 85% > Population in urban areas: 88% > Income group (World Bank): High income


## ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2010

Data refer to litres of pure alcohol per capita (15+).



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2010



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	Average 2003–2005	Average 2008–2010	Change
Recorded	18.0 <sup>†</sup>	12.4 <sup>†</sup>	↘
Unrecorded	1.4	1.4	→
<b>Total</b>	<b>19.4</b>	<b>13.8</b>	↘
Total males / females		19.5   8.2	
WHO European Region	11.9	10.9	

<sup>†</sup>Tourist consumption deducted when at least as many tourists as inhabitants.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2010

Males (15+)	26.6
Females (15+)	13.0
Both sexes (15+)	20.1

Abstainers (%), 2010

	Males	Females	Both sexes
Lifetime abstainers (15+)	8.0	17.1	12.6
Former drinkers* (15+)	18.6	19.6	19.1
Abstainers (15+), past 12 months	26.6	36.7	31.7

\*Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

Prevalence of heavy episodic drinking\* (%), 2010

	Population	Drinkers only
Males (15+)	6.9	9.4
Females (15+)	0.9	1.3
Both sexes (15+)	3.8	5.6

\*Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Patterns of drinking score, 2010

LEAST RISKY < 1 2 3 4 5 > MOST RISKY

## HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2012

	ASDR*		AAF (%)	
Liver cirrhosis, males / females	—	—	—	—
Road traffic accidents, males / females	—	—	—	—

\*Per 100 000 population (15+).

Prevalence of alcohol use disorders and alcohol dependence (%), 2010\*

	Alcohol use disorders**	Alcohol dependence
Males	8.6	5.6
Females	1.9	1.4
Both sexes	5.2	3.4
WHO European Region	7.5	4.0

\*12-month prevalence estimates (15+).

\*\*Including alcohol dependence and harmful use of alcohol.

Years of life lost (YLL) score\*, 2012

LEAST No information available MOST

\*Based on alcohol-attributable years of life lost.

## POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	No / —
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions on on-/off-premise sales of alcoholic beverages: Hours, days / places, density Specific events / intoxicated persons / petrol stations	No, No / Yes, No No / Yes / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.05 / 0.05 / 0.02
Legally binding regulations on alcohol advertising / product placement	No / No
Legally binding regulations on alcohol sponsorship / sales promotion	No / No
Legally required health warning labels on alcohol advertisements / containers	No / No
National government support for community action	Yes
National monitoring system(s)	No