


Egypt

Total population: 78 076 000 > Population aged 15 years and older (15+): 68% > Population in urban areas: 43% > Income group (World Bank): Lower middle income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2010

Data refer to litres of pure alcohol per capita (15+).



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	Average 2003–2005	Average 2008–2010	Change
Recorded	0.2	0.2	→
Unrecorded	0.1	0.2	→
Total	0.3	0.4	→
Total males / females		0.7 / 0.0	
WHO Eastern Mediterranean Region total	0.7	0.7	

Prevalence of heavy episodic drinking* (%), 2010

	Population	Drinkers only
Males (15+)	0.1	0.6
Females (15+)	<0.1	0.2
Both sexes (15+)	<0.1	0.5

*Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2012

	ASDR*		AAF (%)	
	1960	2012	1960	2012
Liver cirrhosis, males / females	122.3	67.8	4.3	8.4
Road traffic accidents, males / females	31.6	8.0	0.1	0.0

*Per 100 000 population (15+).

Years of life lost (YLL) score*, 2012


LEAST < 1 2 3 4 5 > MOST

*Based on alcohol-attributable years of life lost.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Yes (2002/2008) / Yes
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 21 / 21
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 21 / 21
Restrictions for on-/off-premise sales of alcoholic beverages: Hours, days / places, density Specific events / intoxicated persons / petrol stations	No, No / Yes, No No / No / Yes

Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2010



Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2010

Males (15+)	8.6
Females (15+)	0.9
Both sexes (15+)	6.0

Abstainers (%), 2010

	Males	Females	Both sexes
Lifetime abstainers (15+)	86.2	93.1	89.7
Former drinkers* (15+)	5.2	2.6	3.9
Abstainers (15+), past 12 months	91.4	95.7	93.5

*Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

Patterns of drinking score, 2010

LEAST RISKY No information available MOST RISKY

Prevalence of alcohol use disorders and alcohol dependence (%), 2010*

	Alcohol use disorders**	Alcohol dependence
Males	0.4	0.2
Females	<0.1	<0.1
Both sexes	0.2	0.1
WHO Eastern Mediterranean Region	0.3	0.2

*12-month prevalence estimates (15+).

**Including alcohol dependence and harmful use of alcohol.

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	Zero tolerance
Legally binding regulations on alcohol advertising / product placement	Yes / Yes
Legally binding regulations on alcohol sponsorship / sales promotion	Yes / Yes
Legally required health warning labels on alcohol advertisements / containers	Yes / —
National government support for community action	Yes
National monitoring system(s)	No