


# Jordan

Total population: 6 455 000 ► Population aged 15 years and older (15+): 65% ► Population in urban areas: 79% ► Income group (World Bank): Upper middle income


## ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2010

Data refer to litres of pure alcohol per capita (15+).



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2010



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	Average 2003–2005	Average 2008–2010	Change
Recorded	0.5	0.5	→
Unrecorded	0.3	0.2	→
<b>Total</b>	<b>0.8</b>	<b>0.7</b>	<b>→</b>
Total males / females		1.2 / 0.2	
WHO Eastern Mediterranean Region total	0.7	0.7	

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2010

Males (15+)	16.5
Females (15+)	10.9
Both sexes (15+)	15.2

Prevalence of heavy episodic drinking\* (%), 2010

	Population	Drinkers only
Males (15+)	<0.1	0.5
Females (15+)	0.0	0.1
Both sexes (15+)	<0.1	0.4

\*Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2010

	Males	Females	Both sexes
Lifetime abstainers (15+)	91.4	94.3	92.8
Former drinkers* (15+)	1.5	3.4	2.4
Abstainers (15+), past 12 months	92.9	97.8	95.3

\*Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

Patterns of drinking score, 2010

LEAST RISKY No information available MOST RISKY

## HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2012

	ASDR*		AAF (%)	
	20.1	11.2	7.9	13.2
Liver cirrhosis, males / females	46.9	17.8	0.5	0.1

\*Per 100 000 population (15+).

Prevalence of alcohol use disorders and alcohol dependence (%), 2010\*

	Alcohol use disorders**	Alcohol dependence
Males	0.6	0.3
Females	0.1	0.1
Both sexes	0.4	0.2
WHO Eastern Mediterranean Region	0.3	0.2

\*12-month prevalence estimates (15+).

\*\*Including alcohol dependence and harmful use of alcohol.

Years of life lost (YLL) score\*, 2012

LEAST < 1 2 3 4 5 > MOST

\*Based on alcohol-attributable years of life lost.

## POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	No / —	National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.05 / 0.05 / 0.05
Excise tax on beer / wine / spirits	Yes / Yes / Yes	Legally binding regulations on alcohol advertising / product placement	Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18	Legally binding regulations on alcohol sponsorship / sales promotion	Yes / Yes
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18	Legally required health warning labels on alcohol advertisements / containers	No / No
Restrictions for on-/off-premise sales of alcoholic beverages: Hours, days / places, density	Yes, Yes / Yes, Yes	National government support for community action	No
Specific events / intoxicated persons / petrol stations	Yes / Yes / Yes	National monitoring system(s)	No